

PULSATIONS OF THE ELECTRON-POSITRON PLASMA IN THE FIELD OF OPTICAL LASERS

D.B. Blaschke^{1,2,3}, A.V. Prozorkevich⁴, S.A. Smolyansky^{3,4} and A.V. Tarakanov⁴

¹ *Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany*

² *Joint Institute for Nuclear Research, 141980, Dubna, Russia*

³ *Fachbereich Physik, Universität Rostock, Rostock, D-18051, Germany*

⁴ *Physics Department, Saratov State University, 410026, Saratov, Russia*

Abstract

The possibility to observe vacuum electron-positron pair creation due to a powerful optical laser pulse is discussed. We employ a quantum kinetic formulation of the problem with a source term describing the vacuum pair production in a homogeneous electric field with arbitrary time dependence (dynamical Schwinger mechanism). For a periodic field weak in comparison with the critical value $E_{cr} = m^2/|e|$, the electron-positron plasma density changes approximately periodically with twice the field frequency. Under these conditions, the mean value $\langle n \rangle$ for the density per period in the volume λ^3 is a more appropriate characteristic quantity than the residual density n_r , which is taken over an integer number of field periods and calculated using the imaginary time method. The value $\langle n \rangle$ is proportional to the squared field intensity and does not depend on the frequency. We show that in terms of the parameter $\langle n \rangle$ an optical laser can be more effective than a X-ray one. We expect that it is possible to observe the vacuum creation effect not only by means of the planned X-ray free electron lasers but already at present-day optical lasers.

1 INTRODUCTION

QED is considered as the most advanced physical theory, many of its predictions have been proven experimentally with highest available precision. Nevertheless, some questions are discussed till now, e.g., the vacuum pair creation effect by a classical electric field [1]. A complete theoretical description of this effect has been obtained [2, 3, 4, 5], but there is still no experimental proof. The main problem is the high value of the critical electric field strength, necessary to be reached for the pair creation, namely $E_{cr} = 1.3 \times 10^{16} \text{ V/cm}$ for electron-positron pairs. According to the Schwinger formula, the pair creation rate in a constant electric field is

$$\frac{dN}{d^3x dt} = \frac{(eE)^2}{4\pi^3} \sum_{n=1}^{\infty} \frac{1}{n^2} \exp\left(-n\pi \frac{E_{cr}}{E}\right) \quad (1)$$

and therefore exponentially suppressed when $E \ll E_{cr}$. Fortunately, the situation changes qualitatively if the field acts a finite time only [2, 6, 7, 8]. In this case, the Schwinger formula as well as its analog for a monochromatic field (Brezin-Itzykson formula [9]) become inapplicable.

There are a few examples for physical situations where the Schwinger effect can be observed, e.g. relativistic heavy ion collisions [10], neutron stars [11, 12] and focussed laser pulses [13]. The structure of a real laser field is too complicated for the analysis, because the Schwinger effect is non-perturbative and it requires the exact solution of the dynamical equations. That is why the approximation of the homogeneous electric field is used in most cases. According to different estimates [8, 9, 14, 15] the effect of vacuum creation can not be observed with the presently achieved level of laser power, see also [16].

The recent development of laser technology, in particular the invention of the chirped pulse amplification method, has resulted in a huge increase of the light intensity in a laser focal spot [17, 18]. The most advanced lasers produce pulses with intensities reaching 10^{22} W/cm^2 and the pulse duration decreasing down to few oscillation periods. As the construction of X-ray free electron lasers XFELs [19] is now planned, the possibility of the experimental proof of the Schwinger effect attracts attention again. The non-stationary effects become important under conditions of short pulses. We use in our work the kinetic equation approach, which allows us to consider the dynamics of the creation process taking into account the initial conditions [6]. Compared to the other treatments, the approach within the framework of a transport equation contains some new dynamical aspects, such as longitudinal momentum dependence of

the distribution functions and non-markovian character of the time evolution. It takes into account the effects of the field switching and statistics, as well [20]. This approach has been applied already to the periodical field case [7] with near-critical values of the field strength and X-ray frequencies. In particular, it was shown that there is an "accumulation" effect when the intensity of the field is about half critical: the average density of pairs grows steadily with the increase of the field period numbers.

In the present work, we consider the other region of field parameters really achievable nowadays in the optical lasers: $E \ll E_{cr}$ and $\nu \ll m$, where ν is the laser field frequency. We suggest to use in the criterion for the creation efficiency the mean value $\langle n \rangle$ for the density per period in the volume λ^3 is a more appropriate characteristic quantity than the residual density n_r . The latter is taken over an integer number of field periods and calculated using the imaginary time method. The main result is that optical lasers can generate a greater density $\langle n \rangle$ than X-ray ones.

The work is organized as follows. Section 2 contains the statement of the problem and the necessary information about the kinetic equation which is used for the description of vacuum pair creation. We solve this equation numerically for the conditions of the SLAC experiment [21] and study some features of pair production dynamics. We compare here our results obtained on the non-perturbative basis with the predictions of another approach [8] and show that optical lasers can be effective generators of electron-positron pairs during the action of a laser pulse. In Section 3, the low density approximation is considered. It allows to get some analytical results and to make simple estimates. Finally, in Section 4, we discuss some possibilities of direct experimental verification of pair production by high power optical lasers.

2 THE KINETIC EQUATION APPROACH

In the kinetic approach [6], the basic quantity is the distribution function of electrons in the momentum representation $f(\mathbf{p}, t)$. The kinetic equation for this function is derived from the Dirac equation in an external time-dependent field by the canonical Bogoliubov transformation method [2], or by the help of the oscillator representation [22]. This procedure is exact but valid only for the simplest field configurations, e.g., the homogeneous time dependent electric field with the fixed direction

$$\mathbf{E}(t) = (0, 0, E(t)), \quad E(t) = -\dot{A}(t), \quad (2)$$

where the vector potential is given in the Hamiltonian gauge $A^\mu = (0, 0, 0, A(t))$ and the overdot denotes the time derivative. Such a field is not appropriate for a quantitative description of the laser pulse, but can probably be used as qualitative model to estimate results. The corresponding kinetic equation in the collisionless limit has the form [6]

$$\frac{df(\mathbf{p}, t)}{dt} = \frac{1}{2} \Delta(\mathbf{p}, t) \int_{t_0}^t dt' \Delta(\mathbf{p}, t') [1 - 2f(\mathbf{p}, t')] \cos \theta(\mathbf{p}, t', t), \quad (3)$$

where

$$\Delta(\mathbf{p}, t) = eE(t) \frac{\sqrt{m^2 + p_\perp^2}}{\omega^2(\mathbf{p}, t)}, \quad (4)$$

$$\omega(\mathbf{p}, t) = \sqrt{m^2 + p_\perp^2 + [p_\parallel - eA(t)]^2}, \quad (5)$$

$$\theta(\mathbf{p}, t', t) = 2 \int_{t'}^t dt_1 \omega(\mathbf{p}, t_1). \quad (6)$$

Eq. (3) can be transformed to a system of ordinary differential equations, which is convenient for a numerical analysis

$$\begin{aligned} \dot{f} &= \frac{1}{2} \Delta v_1, \\ \dot{v}_1 &= \Delta(1 - 2f) - 2\omega v_2, \\ \dot{v}_2 &= 2\omega v_1, \end{aligned} \quad (7)$$




Figure 1: Time dependence of the density $n(t)$ in the volume λ^3 in the weak periodic field with the parameters $E/E_{cr} = 4.6 \cdot 10^{-6}$ and $\nu/m = 4.29 \cdot 10^{-6}$, corresponding to the SLAC experiment [21] (solid line) and in the near-critical field case of an X-ray laser [19] with $E/E_{cr} = 0.24$ and $\nu/m = 0.0226$ (dashed line).

where v_1, v_2 are real auxiliary functions. The system (7) is integrated via the Runge-Kutta method with the initial conditions $f(\mathbf{p}, t_0) = v_1(\mathbf{p}, t_0) = v_2(\mathbf{p}, t_0) = 0$. The momentum dependence of the distribution function is defined by means of a discretization of the momentum space in a 2-dimensional grid, where the system (7) is solved in each of its nodes. The concrete grid parameters depend on the field strength, where typical values are $\Delta p \approx 0.05 m$ (grid step) and $p_{max} \approx (5 - 10) m$ (grid boundary). The particle number density can be found after that as a moment of the distribution function

$$n(t) = 2 \int \frac{d^3 p}{(2\pi)^3} f(\mathbf{p}, t). \quad (8)$$

Let us consider a harmonic time dependence of the field

$$E(t) = E_m \sin \nu t, \quad A = -\frac{E_m}{\nu} \cos \nu t. \quad (9)$$

The time dependence of density for the field (9) with the parameters $E_m/E_{cr} = 4.6 \cdot 10^{-6}$ and $\nu/m = 4.29 \cdot 10^{-5}$, corresponding to SLAC experiments [21] is shown in Fig. 1 in comparison to the planned X-ray laser [19] with $E/E_{cr} = 0.24$ and $\nu/m = 0.0226$. The pair density oscillates with twice the frequency of the laser field. The density value n_r , which is evaluated in the imaginary time method [8, 15], corresponds to an integer number of field periods, $n_r = n(t = 2\pi/\nu)$, and it is negligible in comparison with the density value n_m corresponding to the electric field maximum, $n_m = n(t = \pi/2\nu)$. The mean density per period $\langle n \rangle$ is of the same order as n_m . For the conditions of the SLAC experiment the ratio of $\langle n \rangle / n_r \approx 3 \cdot 10^{11}$. As a consequence, in spite of the fact that the residual density for the X-ray laser exceeds the one for the optical laser by a large factor, the situation is different regarding the mean density: the optical laser can produce more pairs per volume lambda cubed than the X-ray one.

3 LOW DENSITY APPROXIMATION

The low density approximation $f \ll 1$ can be used in the weak field limit $E \ll E_{cr}$. In that case it is possible to obtain analytic estimates for the residual (n_r) and the maximal (n_m) densities the Eq. (3).




Figure 2: The transverse (left panel) and the longitudinal (right panel) momentum dependence of the distribution function at $t = \pi/2\nu$ (maximum of the field) for the two values of the field strength: $E_m = 4.6 \cdot 10^{-6} E_{cr}$ (solid lines) and $E_m = 3 \cdot 10^{-6} E_{cr}$ (dashed lines) with optical frequency $\nu = 4.26 \cdot 10^{-5} m$.

The particle density in this approximation is

$$n(t) = \frac{e^2}{(2\pi)^3} \int d\mathbf{p} \varepsilon_{\perp}^2 \int_{t_0}^t dt_1 \frac{E(t_1)}{\omega^2(t_1)} \int_{t_0}^{t_1} dt_2 \frac{E(t_2)}{\omega^2(t_2)} \cos \left(2 \int_{t_2}^{t_1} dt_3 \omega(t_3) \right) \quad (10)$$

and can be transformed to [23]

$$n(t) = \frac{1}{2(2\pi)^3} \int d\mathbf{p} \varepsilon_{\perp}^2 \left| \int_{t_0}^t dt_1 \frac{eE(t_1)}{\omega^2(t_1)} \exp \left(2i \int_{t_1}^t dt_3 \omega(t_3) \right) \right|^2. \quad (11)$$

Let us assume additionally that the condition

$$\gamma = \frac{m\nu}{|e|E_m} \gg 1, \quad (12)$$

is satisfied, where γ is the adiabaticity parameter [8]. This relation can be treated as the condition for quasi-classical charge transport in an external field on the time scale $\sim 1/\nu$, if only the pairs are created with vanishing momenta. The latter condition was often used in relation to the longitudinal momentum [24] but the real momentum distribution of the electron-positron pairs has a width of the order of the inverse mass for both transverse and longitudinal momenta, see Fig. 2. The momentum distribution shape varies essentially at the moments of time corresponding to the field minima: a complex quasi-periodic structure with a mean period of about the inverse laser frequency is formed. The mean period value of such a structure decreases proportionally to the number of field periods.

By means of the inequality (12) Eq. (11) is reduced to

$$n(t) = \frac{1}{2(2\pi)^3} \int d\mathbf{p} \frac{\varepsilon_{\perp}^2}{\omega^4} \left| \int_{t_0}^t dt_1 eE(t_1) \exp(2i\omega t_1) \right|^2. \quad (13)$$

The time integral is calculated analytically for the field (9) and $t_0 = 0$ with the result

$$n(t) = \frac{1}{2(2\pi)^3} \int d\mathbf{p} \frac{\varepsilon_{\perp}^2}{\omega^4} \left(\frac{eE_m}{\nu^2 - 4\omega^2} \right)^2 \left\{ \nu^2 (1 + \cos^2 \nu t) + 4\omega^2 \sin^2 \nu t - 2\nu [\nu \cos \nu t \cos 2\omega t + 2\omega \sin \nu t \sin 2\omega t] \right\}. \quad (14)$$

According to Eq. (14) the residual pair density after N periods is $n_r = n(2\pi N/\nu)$ and mean pair density per period $\langle n \rangle$ are estimated as

$$n_r = \frac{1}{(2\pi)^3} \int d^3p \frac{\varepsilon_{\perp}^2}{\omega^4} \left(\frac{eE_m \nu}{\nu^2 - 4\omega^2} \right)^2 \left[1 - \cos \left(\frac{4\pi N \omega}{\nu} \right) \right], \quad (15)$$

$$\langle n \rangle = \frac{1}{4(2\pi)^3} \int d^3p \frac{\varepsilon_{\perp}^2}{\omega^4} \left(\frac{eE_m}{\nu^2 - 4\omega^2} \right)^2 [3\nu^2 + 4\omega^2]. \quad (16)$$

Now we omit the fastly oscillating term in (15) and suppose additionally that

$$\nu \ll m. \quad (17)$$

Then we obtain the simple estimate

$$\frac{\langle n \rangle}{n_r} \sim \left(\frac{m}{\nu} \right)^2. \quad (18)$$

The mean density of electron-positron pairs is defined in this case only by the field amplitude and does not depend on the frequency within a wide range of parameters (12). After the integer period number (when the electric field vanishes) the overwhelming part of pairs is absorbed and the residual density, which is estimated within the usual approach [15], is negligible in comparison with the mean one used above. For the Terawatt Nd-glass laser with the wavelength 527 nm and the field strength $E_m = 6 \cdot 10^{10}$ V/cm [21] we have $m/\nu \approx 2 \cdot 10^5$, so the mean density exceeds the residual one by more than 10 orders of magnitude. According to Fig. 1, there are $\approx 10^5$ pairs in a volume of wavelength cubed on the average for one period of the laser field. The same pair density can be achieved under the conditions of an X-ray laser [7]. Let us notice, that the formula from Ref. [9] for the pair creation probability with the condition (12)

$$w \simeq \frac{(eE)^2}{8} \left(\frac{eE}{2m\omega} \right)^{4m/\omega} \quad (19)$$

gives only a negligible creation probability $\approx 10^{-10^5}$ in this case. This is not surprising because the formula (19) is not applicable for field pulses of finite duration.

4 SUMMARY

The simplest laser field model (9) predicts the existence of a dense electron-positron plasma during a laser pulse duration, which is absorbed almost completely after switching off the field. The mean density is defined by the field strength and does not depend on frequency. The plasma density reaches 10^{18} cm^{-3} within the range of the really achieved fields $10^{10} - 10^{11} \text{ V/cm}$. The usual recipe for an experimental proof of the Schwinger effect suggests that an increase of the residual plasma density can be achieved by an increase of both the frequency and the power of the laser radiation, e.g., via XFEL facilities [19]. It is possible that the simpler way is to try to probe the plasma with some external field in addition to the generating one. SLAC experiments [21] can serve as an example of this type, where a high energy (46 GeV) electron beam crosses the focus of the 527 nm Terawatt pulsed Nd-glass laser and 106 ± 14 positrons above background have been observed. The observed positrons are interpreted as arising from a two-step process: laser photons are backscattered to GeV energies by the electron beam and a collision between the high-energy photon and several laser photons produces an electron-positron pair.

Acknowledgments. This work was supported partly by Russian Federations State Committee for Higher Education under grant E02-3.3-210 and Russian Fund of Basic Research (RFBR) under grant 03-02-16877. S.A.S. acknowledges support by the DFG Graduiertenkolleg 567 "Stark korrelierte Vielteilchensysteme" at the University of Rostock and by DFG grant No. 436 RUS 117/78/04.

References

- [1] F. Sauter, Z. Phys. **69**, 742 (1931); W. Heisenberg and H. Euler, Z. Phys. **98**, 714 (1936); J. Schwinger, Phys. Rev. **82**, 664, (1951).

- [2] A.A. Grib, S.G. Mamaev, and V.M. Mostepanenko, *Vacuum Quantum Effects in Strong External Fields* (Friedmann Lab. Publ., St.-Petersburg, 1994).
- [3] A.I. Nikishov, Tr. Fiz. Inst. Akad. Nauk SSSR, **111**, 152 (1979).
- [4] W. Greiner, B. Müller, and J. Rafelski, *Quantum Electrodynamics of Strong Fields*, (Springer, Berlin, 1985).
- [5] E.S. Fradkin, D.M. Gitman, and S.M. Shvartsman, *Quantum Electrodynamics with Unstable Vacuum*, (Springer-Verlag, Berlin, 1991).
- [6] S.M. Schmidt, D. Blaschke, G. Röpke, S.A. Smolyansky, A.V. Prozorkevich, and V.D. Toneev, Int. J. Mod. Phys. E, **7**, 709 (1998).
- [7] R. Alkofer, M.B. Hecht, C.D. Roberts, S.M. Schmidt, and D.V. Vinnik, Phys. Rev. Lett. **87**, 193902, (2001); C.D. Roberts, S.M. Schmidt, and D.V. Vinnik, *ibid*, **89**, 153901, (2002).
- [8] S.V. Popov, JETP Lett. **74**, 133, (2001); Phys. Lett. A **298**, 83 (2002).
- [9] E. Brezin and C. Itzykson, Phys. Rev. D **2**, 1191, (1970).
- [10] A. Casher, H. Neuberger, and A. Nussinov, Phys. Rev. D **20**, 179 (1979).
- [11] V.S. Beskin, A.V. Gurevich, and Ya.N. Istomin, *Physics of the Pulsar Magnetosphere* (Cambridge: Cambridge Univ. Press, 1993).
- [12] R. Ruffini, L. Vitagliano, and S.-S. Xue, Phys. Lett. B **559**, 12 (2003).
- [13] B. Richards and E. Wolf, Proc. Roy. Soc. A (London) **253**, 358 (1959).
- [14] F.V. Bunkin and I.I. Tugov, Dokl. Akad. Nauk. SSSR **187**, 541 (1964) [Sov. Phys. Dokl. **14**, 678 (1969)].
- [15] M.S. Marinov and V.S. Popov, Fortschr. Phys. **25**, 373 (1977).
- [16] S.S. Bulanov, Phys. Rev. E **69**, 0326408 (2004); S.S. Bulanov, N.B. Narozhny, and V.D. Mur, V.S. Popov, hep-ph/0403163; S.S. Bulanov, A.M. Fedotov, and F. Pedoraro, hep-ph/0409301.
- [17] G.A. Mourou, C.P.J. Barty, and M.D. Perry, Phys. Today **51**, No.1, 22 (1998).
- [18] S.V. Bulanov, T. Esirkepov, and T. Tajima, Phys. Rev. Lett. **91**, 085001 (2003), *ibid*. **92**, 159901 (2004)(E).
- [19] A. Ringwald, Phys. Lett. B **510**, 107 (2001).
- [20] S.M. Schmidt, D. Blaschke, G. Röpke, A.V. Prozorkevich, S.A. Smolyansky, and V.D. Toneev, Phys. Rev. D, **59**, 094005 (1999); J. C. Bloch, V. A. Mizerny, A.V. Prozorkevich, C. D. Roberts, S.M. Schmidt, S.A. Smolyansky, and D. V. Vinnik, Phys. Rev. D, **60**, 1160011 (1999).
- [21] C. Bula et al., Phys. Rev. Lett. **76**, 3116 (1996); D.L. Burke et al., Phys. Rev. Lett. **79**, 1626 (1997).
- [22] V.N. Pervushin, V.V. Skokov, A.V. Reichel, S.A. Smolyansky, and A.V. Prozorkevich, hep-th/0307200.
- [23] A. V. Prozorkevich, A. Reichel, S. A. Smolyansky, A. V. Tarakanov, Proceedings of SPIE, **5476**, 68 (2004)
- [24] A. Casher, H. Neuberger, and S. Nussinov, Phys. Rev. D, **20**, 179 (1979).